

Supplemental Table 1 Baseline characteristics of the patients by those who changed facility during the treatment and those who did not

Variable		Moved centers during the treatment 173,025 (66%)	Did not move during treatments 89,184 (34%)	P
Age at diagnosis, Median (range)		70 (40-90)	71 (40-90)	
Sex	Male	86,856 (50.2%)	44,612 (50.0%)	0.39
	Female	86,169 (49.8%)	44,572 (50.0%)	
Race	White	144,676 (84.7%)	72,609 (82.2%)	0.001
	Black	20,694 (12.1%)	12,224 (13.8%)	
	neither White nor Black	5,474 (3.20%)	3,510 (4.0%)	
Education	>=13% NHD	74,864 (43.5%)	38,657 (43.4%)	0.7
	<13% NHD	97,241 (56.5%)	50,362 (56.6%)	
Income	>=\$35,000	100,172 (58.2%)	51,698 (58.1%)	0.5
	<35,000	71,833 (41.8%)	37,281 (41.9%)	
Place of Living	Urban	165,225 (98.1%)	85,387 (98.1%)	0.6
	Rural	3,205 (1.9%)	1,628 (1.9%)	
Insurance Status	Yes	163,612 (97.09%)	85,051 (96.83%)	0.001
	No	4,899 (2.91%)	2,785 (3.17%)	
Charlson/Deyo Score	0	115,217 (66.6%)	55,260 (62.0%)	0.001

	1	41,484 (24.0%)	22,484 (25.2%)	
	>=2	16,324 (9.4%)	11,440 (12.8%)	
M stage	M0	85,028 (50.5%)	31,471 (37.2%)	0.001
	M1	83,287 (49.5%)	53,209 (62.8%)	
Chemotherapy	Yes	88,869 (51.4%)	41,684 (46.7%)	0.001
	No	84,156 (48.6%)	47,500 (53.3%)	
Radiation Therapy	Yes	26,333 (15.2%)	8,741 (9.8%)	0.001
	No	146,692 (84.8%)	80,443 (90.2%)	
Immunotherapy	Yes	612 (0.4%)	286 (0.3%)	0.2
	No	172,413 (99.6%)	88,898 (99.7%)	
Year of Diagnosis	2004-2010	115,561 (66.8%)	10,204 (11.4%)	0.001
	2011-2016	57,464 (33.2%)	78,980 (88.6%)	